

Collision-Dependent Atom Tunneling Rate in Bose-Einstein Condensates

B. R. da Cunha and M. C. de Oliveira

Instituto de Física Gleb Wataghin, Universidade Estadual de Campinas, 13083-970, Campinas - SP, Brazil.

(Dated: February 2, 2008)

We show that the interaction (cross-collision) between atoms trapped in distinct sites of a double-well potential can significantly increase the atom tunneling rate for special trap configurations leading to an effective linear Rabi regime of population oscillation between the trap wells. The inclusion of cross-collisional effects significantly extends the validity of the two-mode model approach allowing it to be alternatively employed to explain the recently observed increase of tunneling rates due to nonlinear interactions.

PACS numbers: 03.75.Lm, 74.50.+r

Atom optics and the physics of ultracold matter waves have witnessed rapid theoretical and experimental progress since the achievement of atomic vapor Bose-Einstein condensates (BECs) [1]. The interest in such a system is quite wide ranged as it has opened new technological frontiers [2, 3], and has renewed the investigation on many-body physics fundamental issues once BECs trapped in optical lattices offers a powerful toolbox for implementation of condensed matter model systems with highly controllable parameters [4, 5, 6].

The simplest system to exhibit non-trivial many-body phenomena is the two-mode BEC trapped in a double well potential. For this system it was previously predicted [7, 8] a *plethora* of dynamical regimes, including Josephson oscillation [9] of atoms between distinct trap sites, and macroscopic self-trapping of atomic populations, where coherent tunneling is suppressed when the number of trapped atoms exceeds a critical value. The first realization of an atomic single BEC Josephson junction has indeed experimentally demonstrated [10] those predictions [7, 8, 9]. Moreover an increase of one order of magnitude of the atomic tunneling rate (from 2 to 25 Hz) due to nonlinear atomic interactions was then observed, which cannot be explained inside those previous models. In those models [7, 8] a two-mode approximation (TMA) is invariably applied, in which each trap well is populated by one single localized condensate mode. However due to a localization assumption the BEC dynamics is considered by regarding only self-collisions (on-site interaction) between atoms of individual condensate modes. Notwithstanding, distinct trap configurations can be as such that cross-collisions (off-site interaction) of atoms in distinct BEC modes cannot be neglected and the observed coherent phenomena are affected by cross-collision induced effects. This situation has been previously considered either by directly integrating a non-polynomial Schrödinger equation [10, 11] or by extending the TMA through Gross-Pitaevskii equations [12].

In this paper we investigate the cross-collisional effects over the quantum dynamics of a BEC trapped in a double-well potential inside the TMA. By appropriate manipulation of the trapping potential the sys-

tem shows a very rich quantum dynamics related to the cross-collisional rates. In absence of cross-collisions, self-trapping is observed in agreement with previous theoretical predictions [7, 8]. Remarkably, the inclusion of cross-collisional effects can inhibit self-trapping, resulting in an effective coherent oscillation (effective Rabi regime) of atoms between the two wells with an effective Rabi frequency, dependent on the cross-collision rate and the number of atoms in the sample. Moreover cross-collisions increase the range of validity of the TMA (when the many-body interactions produce only slight changes on the ground-state properties of the individual potential sites). The new limit can be as such that the TMA is indeed valid for the number of atoms present in the experiment of Ref. [10], and thus can be employed for alternative explanation [10, 12] of the observed increase of tunnelling rate through off-site interactions.

We consider an atomic BEC trapped in a symmetric double-well potential $V(\mathbf{r})$ with minima at $\mathbf{r}_1$, and $\mathbf{r}_2$, such that $V(\mathbf{r}_{1,2}) = 0$, the many-body Hamiltonian is given by

$$\hat{H} = \int d^3r \hat{\psi}^\dagger(\mathbf{r}) \left(-\frac{\hbar^2}{2m} \nabla^2 + V(\mathbf{r}) \right) \hat{\psi}(\mathbf{r}) + \frac{1}{2} U_0 \int d^3r \hat{\psi}^\dagger(\mathbf{r}) \hat{\psi}^\dagger(\mathbf{r}) \hat{\psi}(\mathbf{r}) \hat{\psi}(\mathbf{r}), \quad (1)$$

where m is the atomic mass, $U_0 = 4\pi\hbar^2 a/m$ measures the strength of the two-body interaction, a is the s-wave scattering length, $\hat{\psi}(\mathbf{r}, t)$ and $\hat{\psi}^\dagger(\mathbf{r}, t)$ are the Heisenberg picture field operators. We must emphasize that throughout our calculations we assume that the barrier height separating the two wells is sufficiently larger than the condensate chemical potential, $V_0 > \mu_c$, so that a TMA can be considered to describe the system [13]. The derivation of the two-mode Hamiltonian in a double-well trap follows the standard procedures of Ref. [7] - The lower energy eigenstates of the global double-well are approximated as the symmetric and anti-symmetric combinations $u_\pm(\mathbf{r}) \approx \frac{1}{\sqrt{2}}[u_1(\mathbf{r}) \pm u_2(\mathbf{r})]$, and the field operators are expanded in terms of the local modes $u_j(\mathbf{r})$, $j = 1, 2$ and the Heisenberg picture annihilation operator as $\hat{a}_j(t) = \int d^3r u_j^*(\mathbf{r}) \hat{\psi}(\mathbf{r}, t)$ so that

$[\hat{a}_j, \hat{a}_k^\dagger] \simeq \delta_{jk}$. The potential expanded around each minimum is $V(\mathbf{r}) = \tilde{V}^{(2)}(\mathbf{r} - \mathbf{r}_j) + \dots$ $j=1,2$, where $\tilde{V}^{(2)}(\mathbf{r} - \mathbf{r}_j)$ is the harmonic approximation to the potential in the vicinity of each minimum. The normalized single-particle ground-state $u_j(\mathbf{r})$ of the local potential $\tilde{V}^{(2)}(\mathbf{r} - \mathbf{r}_j)$, with energy E_0 , defines the local mode solutions of the individual wells. The tunneling frequency Ω between the two minima is then given by the energy level splitting of these two lowest states, $\Omega = 2R/\hbar$, where $R = \int d^3\mathbf{r} u_1^*(\mathbf{r})[V(r) - \tilde{V}^{(2)}(\mathbf{r} - \mathbf{r}_1)]u_2(\mathbf{r})$. By noticing that the total number of atoms N is a conserved quantity, after some algebraic manipulation, the following Hamiltonian is then obtained in the interaction picture

$$\hat{H} = \hbar[2\Lambda(N-1) + \Omega](\hat{a}_1^\dagger \hat{a}_2 + \hat{a}_2^\dagger \hat{a}_1) + \hbar\eta(\hat{a}_1^\dagger \hat{a}_2 + \hat{a}_2^\dagger \hat{a}_1)^2 + \hbar(\kappa - \eta)[(\hat{a}_1^\dagger)^2(\hat{a}_1^2) + (\hat{a}_2^\dagger)^2(\hat{a}_2^2)] + \hbar\eta N(N-2) \quad (2)$$

where $\kappa = \frac{U_0}{2\hbar V_{eff}}$ is the self-collision rate and $\eta = (\frac{U_0}{2\hbar}) \int d^3\mathbf{r} u_j^* u_j u_i^* u_{0j}$, and $\Lambda = (\frac{U_0}{2\hbar}) \int d^3\mathbf{r} u_j^* u_j u_i^* u_i$, are cross-collisional rates. Here $V_{eff}^{-1} = \int d^3\mathbf{r} |u_j|^4$ is the j -mode effective volume. It is immediate that for $\hbar\kappa \rightarrow 0$, the Hamiltonian (2) reduces to that previously investigated in the study of tunneling in condensates [9]. Also for $\hbar\eta \rightarrow 0$ and $\hbar\Lambda \rightarrow 0$, the Hamiltonian (2) reduces itself to that discussed in [7, 8], accounting for tunneling oscillations as well as for population self-trapping. As $\eta \rightarrow \kappa$ in Hamiltonian (2) a new dynamical regime of stable long-time tunneling (hereafter called effective Rabi regime) can be attained.

Introducing the Schwinger angular momentum operators $\hat{J}_x = (\hat{a}_2^\dagger \hat{a}_2 - \hat{a}_1^\dagger \hat{a}_1)/2$, $\hat{J}_y = i(\hat{a}_2^\dagger \hat{a}_1 - \hat{a}_1^\dagger \hat{a}_2)/2$, $\hat{J}_z = (\hat{a}_1^\dagger \hat{a}_2 + \hat{a}_2^\dagger \hat{a}_1)/2$, the Hamiltonian (2) can be rewritten as

$$\hat{H} = \hbar\Omega' \hat{J}_z + 4\hbar\eta \hat{J}_z^2 + 2\hbar(\kappa - \eta) \hat{J}_x^2, \quad (3)$$

where we have neglected constant energy shifts dependent on N and have defined a new effective tunneling rate $\Omega' \equiv 2[\Omega + 2\Lambda(N-1)]$, which is dependent on the number of atoms in the atomic sample and the cross-collisional rate Λ . In fact Λ is very small compared to Ω , but for a sufficiently large number of atoms the additional tunneling term may lead to observable effects whenever $2\Lambda(N-1) \geq \Omega$. Furthermore the third term of Eq. (3) shows that the cross-collisional rate η competes with the self-collision rate κ leading to an effective on-site collision rate $\kappa' \equiv \kappa - \eta$. Since $0 \leq \eta \leq \kappa$, κ' on Eq. (3) can be disregarded in a trap configuration where $\eta \rightarrow \kappa$ with $\Omega' \gg \kappa'$. To infer the validity and implications of such a regime we consider a specific trapping potential [14] of the form $V(r) = b(x^2 - \frac{d}{2b})^2 + \frac{1}{2}m\omega_t^2(y^2 + z^2)$, where the inter-well coupling occurs along x , and ω_t is the trap frequency in the y - z plane. This potential has fixed points at $q_0^2 = \frac{d}{2b}$. The position uncertainty for a harmonic oscillator in the ground state is $x_0 \equiv \sqrt{\frac{\hbar}{2m\omega_0}}$, with $\omega_0 = \sqrt{4d/m}$. For a suitable choice of the barrier height

only two energy eigenstates lie beneath the barrier. Assuming $\omega_t = \omega_0$ the local mode on each well is then given by $u_j(r) = (\frac{1}{2\pi x_0^2})^{\frac{3}{4}} \exp\left(\frac{-[(x_j - q_0)^2 + y_j^2 + z_j^2]}{4x_0^2}\right)$. The collision rates may then be evaluated to give $\kappa = (\frac{U_0}{16\hbar})(\frac{1}{\pi x_0^2})^{3/2}$, $\eta = \kappa \exp(-q_0^2/2x_0^2)$, and $\Lambda = \kappa \exp(-3q_0^2/4x_0^2)$, while $\Omega = \frac{q_0^2 \omega_0}{x_0^2} \exp(-q_0^2/2x_0^2)$.

Firstly we shall investigate the validity of the localization of the TMA considered in the present description. It is valid only if the many-body interactions produce small modifications on the ground-state of the individual potential wells [15]. In the absence of cross-collisions [7] the TMA is limited to $\hbar\omega_0 = \frac{\hbar^2}{2mx_0^2} \gg N\hbar\kappa = N\frac{\hbar|U_0|}{2\hbar V_{eff}}$ and thus the BEC is limited to a few 100 atoms. The inclusion of cross-collision necessarily increases the effective mode volume, which in terms of the effective on-site collision rate κ' is given by $V'_{eff} \equiv V_{eff}/(1 - e^{-q_0^2/2x_0^2})$. For the trapping potential considered, $V_{eff} \sim 8\pi^{3/2}x_0^3$, and the TMA is valid whenever

$$N \ll \frac{2\pi^{1/2}x_0}{|a|(1 - e^{-q_0^2/2x_0^2})}. \quad (4)$$

For suitable values of the trap parameters, it is thus possible to have very large numbers at the right of the inequality above. Hence the considered TMA, with cross-collision included, can be appropriately employed for the description of the observed phenomena in present experimental setups [10] for numbers of atoms $N \gtrsim 10^3$.

Typically $\eta, \Lambda \ll \Omega$, but cross-collisional effects can only be neglected if $2\Lambda(N-1)/\Omega \ll 1$, *i. e.* if $(N-1)\frac{a_s x_0}{\pi^{1/2} q_0} e^{-q_0^2/4x_0^2} \ll 1$. By considering the experimental values from Ref. [10] for $N = 1300$, $a_s = 5.3$ nm for ^{87}Rb atoms, and $q_0 = 2.2\mu\text{m}$, we obtain that only for $x_0 \ll 0.82 q_0$ is that cross-collisional effects can be neglected. This bound value however depends strongly on the number of atoms in the sample and decreases rapidly as N is increased. For example, if the number of atoms is increased in one order of magnitude to $N = 13000$ then the cross-collisional effects can be neglected only if $x_0 \ll 0.366 q_0$. Furthermore by equating Ω' to the observed frequency of oscillation of 25 Hz in the experiment performed by Albiez *et al.* [10] and Ω to the expected frequency of tunneling of 2 Hz we find with the above parameters that only for $x_0 \approx 6.56 q_0$ is that the tunneling frequency would significantly be increased due to non-linear off-site interactions. Of course, inside the applied TMA this situation corresponds to that of two strongly overlapped modes, which would compromise the validity of the localization assumption and the considered gaussian eigenfunctions. However the relation between x_0 and q_0 is not linear and depends strongly on N and on q_0 as well. For a fixed q_0 the x_0 to q_0 ratio decreases rapidly as N is increased. Again, for $N = 13000$ the substantial increase in the tunneling frequency would be observed for $x_0 \approx 0.89 q_0$. We remark that the above analysis could

be similarly applied by considering the local modes as approximately given by the excited states of the harmonic oscillator. In such a case the localization conditions in relation to the distance between minima of the trapping potential would be significantly relaxed [18].




FIG. 1: Numerical results for the eigenvalues of Eq. 3 in the eigenbasis of $\hat{J}_x$. The energy spectra is shown for different regimes of the η/κ ratio. Each spectrum has been displaced by a constant factor.

We now turn to the description of the dynamical regimes attained by the Hamiltonian (3). In Fig. 1 we plot the numerically calculated eigenvalues spectra for a condensate with 1000 atoms and fixed $\Omega/\kappa = 50$ ratio. Each curve in the plot represents a specific η/κ ratio as depicted in the figure. For $\eta/\kappa = 1/1000$ it is immediately verified that the eigenvalues correspond to those of the model without cross-collision discussed in Ref. [7]. The low-excited eigenstates are close to the eigenstates of the $\hat{J}_z$ operator given by $|j, -j\rangle_z$. Highly excited eigenstates are closer to the eigenstates of the operator $\hat{J}_x^2$, given approximately by $|j, 0\rangle_x$. This last state suggests an ideal situation for occurrence of population self-trapping for large N . The inflexion point for intermediate eigenvalues represents the frontier for the occurrence of self-trapping. As the ratio κ/η is increased the eigenvalues behavior changes dramatically due to the influence of the term proportional to $\hat{J}_x^2$. An outstanding feature is that the inflexion point is shifted to higher eigenvalues for $\kappa/\eta = 1/100 - 1/10$ and completely disappears for $\kappa/\eta \geq 1/5$. For $\kappa/\eta \approx 1$ the term proportional to $\hat{J}_x^2$ is negligible in comparison to the other terms and the Hamiltonian eigenstates get as closer to the $\hat{J}_z$ eigenstates as $\kappa/\eta \rightarrow 1$.

The equations of motion for the angular momentum operators are given by

$$\dot{\hat{J}}_x = -\hbar\Omega'\hat{J}_y - 4\hbar\eta[\hat{J}_y, \hat{J}_z]_+ \quad (5)$$

$$\dot{\hat{J}}_y = \hbar\Omega'\hat{J}_x - 2\hbar(\kappa - 3\eta)[\hat{J}_z, \hat{J}_x]_+ \quad (6)$$

$$\dot{\hat{J}}_z = 2\hbar(\kappa - \eta)[\hat{J}_y, \hat{J}_x]_+ \quad (7)$$

One can immediately see that if the cross-collision term

is strong enough so that $\kappa - \eta \ll \Omega'$ then

$$\dot{\hat{J}}_z \approx 0 \quad (8)$$

and thus $\hat{J}_z$ being a constant of motion the set of Eqs. (5, 6, and 8) is solved for $\hat{J}_x$ as

$$\hat{J}_x(t) = \hat{J}_x(0) \cos \Omega'' t + \hat{J}_y(0) \sin \Omega'' t, \quad (9)$$

where $\Omega'' \equiv \Omega' + 8\eta J_z = [\Omega + 2\Lambda(N-1)] + 8\eta J_z$ is the new Rabi frequency, which now depends on both the number of atoms and the initial condition for J_z through the nonlinear cross-collision rates Λ and η , respectively. Since $\eta \ll \Omega$, J_z value only gives a small shift in Ω'' . The effective Rabi regime of $\eta \rightarrow \kappa$ is justified whenever $\kappa - \eta \ll \Omega''$.

In Fig. 2 the time evolution of the mean value $\langle \hat{J}_x \rangle$ as given by the general system of Eqs. (5-7) for distinct η/κ ratio. We suppose an initial state localized in one well as $|j, -j\rangle_x$. Similarly to Ref. [7] due to intrinsic quantum fluctuations in the initial condition there are some oscillations of the quantum mean decay [19]. In Fig. 2(a) the revival of the oscillation that occurs at later times is again due to the discrete spectrum of the many-body Hamiltonian [20, 21]. In Fig. 2(b) the self-trapping of the atomic population is shown. This result should be compared to that of Fig. 2 from Ref. [10]. More interesting though are the new features in the quantum dynamics due entirely to the presence of cross-collisions appearing in Fig. 2(c). Notice that even when cross-collisions are relatively small ($\kappa/\eta \sim 10$) coherent tunneling dominates, suppressing the initial self-trapping. As the cross-collisions become stronger the collapse and revival dynamics changes. For large intervals of time there is a total collapse of the mean value, which lasts for large periods of time in such a way that this period decreases as η gets closer to κ approaching the limit where it tends to zero when the effective Rabi regime takes place and there is no collapse and revival. Notice that the decreasing amplitude modulation in the collapse and revival is due to nonlinearities introduced in Eqs. (5-7) by cross-collisions. As $\eta \rightarrow \kappa$ however, the amplitude of coherent oscillation of the population difference goes to a constant value with oscillating frequency Ω'' .

To push the discussion presented here further, it is interesting to consider the impact of the cross-collisional terms in related many wells systems [6]. The dynamics is richer and additional effects may occur as consequence of the overlap of neighboring condensed modes. The results found for the double-well case can be extended and set into the more general framework of hopping in many wells systems. It is possible to render a general discussion of the cross-collisional terms for the derivation of a discrete extended Bose-Hubbard model [18] and of the Wannier-Stark effect [22] based solely on ground state trap solutions. For instance, a n -mode *ansatz* of the natural

extension of the Hamiltonian in Eq.(2) results in a n -mode extended Bose-Hubbard Hamiltonian

$$\begin{aligned}\hat{H} = & \sum_l (\kappa_l - \eta_{l,l+1}) \hat{n}_l (\hat{n}_l - 1) \\ & + \sum_l \eta_{l,l+1} (\hat{a}_{l+1}^\dagger \hat{a}_l + H.c)^2 \\ & + \sum_l [\Omega_{l,l+1} + 2\Lambda_{l,l+1}(2\hat{n}_l - 1)] (\hat{a}_{l+1}^\dagger \hat{a}_l + H.c),\end{aligned}\quad (10)$$

with all parameters related to the those for the double-well system. The dynamical regimes given by (10) have in part a similar aspect to the ones presented for the double-well potential but can be extended since all the on-site and off-site parameters can be independently varied. Those issues are being further investigated in the context of phase transitions and shall be addressed elsewhere [23].




FIG. 2: Numerical results for the mean value of the population imbalance $\langle \hat{J}_x \rangle$ for $N=100$, $\kappa N = 2.0$, (a) $\eta = \kappa/10$, (b) $\eta = \kappa/100$ and (c) $\eta = \kappa/2$. For convenience we normalize the time in units of the single particle tunneling period so that in these units $\Omega = 1$. Time evolution found by integrating Schrödinger equation in the $\hat{J}_x$ basis for the Hamiltonian in Eq. 3.

In summary we employed the TMA to model an atomic two-mode BEC trapped in a double well potential considering cross-collisions between the condensed modes. Cross-collisions strongly inhibit the self-trapping phenomenon even for small values of the cross-collision rate η . For a given trap geometry the eigenvalues of the many-body Hamiltonian determine a new dynamics resulting in an effective coherent oscillation (effective Rabi regime) of the population in both wells. In the limit $\eta \rightarrow \kappa$ for $\kappa - \eta \ll \Omega$ an effective Rabi regime takes place with an effective Rabi frequency, Ω , which is explicitly dependent on the total number of atoms. The limit of valid-

ity of such a two-mode approximation increases within the assumption of strong cross-collisions in such a way that it may be in accordance with experimental data [10] observed for the increase of tunneling frequency due to nonlinear interaction. As a last result the collapse and revival dynamics is changed resulting in large periods of total collapse. As $\eta \rightarrow \kappa$, however the collapse and revival frequency increases attaining the regime of Rabi population oscillation for $\eta \sim \kappa$. It is remarkable that similar results to those previously given in exact numerical calculations [10, 11, 12] could be derived inside the TMA just by adding cross-collisional effects. We hope that the above considerations bring some contribution to experimental investigation.

The authors would like to acknowledge partial financial support from FAPESP under project #04/14605-2, from CNPq and from FAEPEX-UNICAMP.

-
- [1] M.H. Anderson *et al.*, Science **269**, 198 (1995); K.B. Davis *et al.*, Phys. Rev. Lett. **75**, 3969 (1995).
 - [2] D. Jaksch *et al.*, Phys. Rev. Lett. **85**, 2208 (2000).
 - [3] M.C. de Oliveira, Phys. Rev. A **67**, 022307 (2003).
 - [4] A. J. Leggett, Rev. of Mod. Phys. **73**, 307 (2001).
 - [5] M. Greiner *et al.*, Nature **415**, 39 (2002).
 - [6] D. Jaksch and P. Zoller, Annals of Physics (N.Y.) **315**, 52 (2005), and references therein.
 - [7] G.J. Milburn *et al.*, Phys. Rev. A **55**, 4318 (1997); J. F. Corney and G.J. Milburn, Phys. Rev. A **58**, 2399 (1998).
 - [8] S. Raghavan *et al.*, Phys. Rev. A **59**, 620 (1999); S. Raghavan, A. Smerzi, and V. M. Kenkre, Phys. Rev. A **60**, 1787 (1999); S. Giovanazzi, A. Smerzi, and S. Fantoni, Phys. Rev. Lett. **84**, 4521 (2000).
 - [9] J. Javanainen, Phys. Rev. Lett. **57**, 3164 (1986).
 - [10] M. Albiez *et al.*, Phys. Rev. Lett. **95**, 010402 (2005).
 - [11] L. Salasnich, A. Parola, and L. Reatto, Phys. Rev. A **65**, 043614 (2002).
 - [12] D. Ananikian and T. Bergeman, Phys. Rev. A **73**, 013604 (2006).
 - [13] J. E. Williams, Phys. Rev. A **64**, 013610 (2001).
 - [14] We remark that the results obtained are qualitatively equivalent for any double-well trapping potential.
 - [15] Otherwise the Thomas-Fermi approximation [16, 17] should be applied instead of the TMA.
 - [16] G. Baym and C. J. Pethick, Phys. Rev. Lett. **76**, 6 (1996).
 - [17] M. Edwards *et al.*, Phys. Rev. A **53**, 1950 (1996).
 - [18] V. W. Scarola and S. Das Sarma, Phys. Rev. Lett. **95**, 033003 (2005).
 - [19] P. W. Anderson, *Basic Notions of Condensed Matter Physics* (addison-Wesley, Reading, MA, 1984).
 - [20] L. Bernstein, J. C. Eilbeck, and A. C. Scott, Nonlinearity **3**, 293 (1990).
 - [21] A. Cheffles and S. M. Barnett, J. Mod. Opt. **43**, 709 (1996).
 - [22] A. Buchleitner and A. R. Kolovsky Phys. Rev. Lett. **91**, 253002 (2003).
 - [23] B. R. da Cunha, R. C. Farias, and M. C. de Oliveira, *unpublished*.